

The perverse timing of stepping the equation counter

May 23, 2018

1 Equations

Note how equations increment the equation counter at the *beginning*, even if later on, the equation tag is removed or set explicitly!

Normal (w/eq=0)	$N = 1$	(1)
-----------------	---------	-----

Nonumber before (w/eq=1)	$N = 1$	
--------------------------	---------	--

Nonumber after (w/eq=1)	$N = 2$	
-------------------------	---------	--

Numbered, Tag before (w/eq=1)	$N = 1$	(1')
-------------------------------	---------	------

Numbered, Tag after (w/eq=1)	$N = 2$	(1'')
------------------------------	---------	-------

Other variations:	$N = 2$	(1 primed)
	$N = 2$	(1 in text)

UnNumbered (w/eq=1)	$N = 1$	
---------------------	---------	--

UnNumbered, Tag before (w/eq=1)	$N = 1$	(1''')
---------------------------------	---------	--------

UnNumbered, Tag after (w/eq=1)	$N = 1$	(1''')
--------------------------------	---------	--------

Now (w/eq=1).

2 Eqnarrays

Note how eqnarrays *also* increment at the beginning! *Even* eqnarray*!

Mostly numbered, unnumbered before and after (w/eq=1)

$$N = 2 \tag{2}$$

$$N = 3 \tag{3}$$

$$N = 4$$

$$N = 4$$

$$N = 4 \tag{4}$$

Unnumbered (w/eq=4)

$$N = 5$$

$$N = 5$$

Now (w/eq=4).

3 AMS alignments

However, note how AMS alignments *defer* incrementing the equation counter until the end of line mark!

Align numbered, some unnumbered before, after, some tagged before, after (w/eq=4)

$$N = 4 \tag{5}$$

$$N = 5 \tag{6}$$

$$N = 6$$

$$N = 6$$

$$N = 6 \tag{7}$$

$$N = 7 \tag{7'}$$

$$N = 7 \tag{7''}$$

Align numbered, some tagged before, after (w/eq=7)

$$N = 7$$

$$N = 7 \tag{7'''} \tag{7''''}$$

$$N = 7 \tag{7''''}$$

$$N = 7$$

Now (w/eq=7).